



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

morants destroy salmon, nor did stomach examinations. Of 32 stomachs examined, 5 were empty, 3 contained unrecognizable food, 16 contained sculpins, 5 herrings, one capelin, one eel, and 2 tomcod or allied fishes.

Furthermore, the bulk of evidence shows that salmon have lately been increasing from year to year. "The cormorants are also generally increasing in number, the rookeries are enlarging and new ones being established. These facts taken together do not indicate that the cormorants are markedly harmful to the salmon."*—W. L. M.

* Mus. Bul. 13, Canada Dept. of Mines, 1915, p. 14.

INCUBATION PERIOD OF THE GANNET.

In the very interesting list of periods of incubation, as noted in different birds, contributed to your pages by Mr. F. L. Burns, the period allowed for the Gannet (*Sula bassana* (L.)) is given at 39 days, but this seems too short.

An egg laid on April 22d at the Bass Rock on the east coast of Scotland, and at once inscribed with the date by Mr. J. M. Campbell, the lighthouse keeper, was not hatched until June 5th, which gives a period of 44 days.

Another Gannet's egg, laid in confinement at Brighton in Sussex, is recorded by the late Mr. E. T. Booth to have hatched out about the 43d day, as stated in "The Gannet" (p. 355), where the *pros* and *cons* of the subject are discussed at some length and different opinions quoted.

J. H. GURNEY.

Keswick Hall, Norwich.

A NEW LOUISIANA BIRD.

The observance of a solitary species of bird life has not only supplied the Louisiana list with a new bird but with the one order of the seventeen found in the United States that has been missing since the avian life of the state has been studied by those who preceded Audubon and those who have followed him.

The species observed was a Wilson petrel (*Oceanites oceanicus*), sometimes known as one of "Mother Cary's Chickens," of the order *Tubinares*, or Tube-nosed swimmers, which includes, besides the petrels, the fulmars and shearwaters.

The discovery was made by Herbert K. Job, head of the Department of Applied Ornithology of the National Association of Audubon Societies, and Stanley Clisby Arthur, ornithologist of the Conservation Commission of Louisiana, while they were on an expe-